

Gonyaulax scrippsae Kofoid

Univ. Calif. Publ. Zool. 8: 228, pl. 13, figs 26, 27; pl. 16, fig. 38 (1911)

Type locality: U.S.A. (California)

Australian distribution: Qld (Gulf of Carpentaria; cf.; cysts), N.S.W. (Sydney Harbour, Port Hacking, off Sydney), W.A. (Swan River)

Extra-Australian distribution: U.S.A. (California), Atlantic Ocean, British Isles, Mediterranean Sea, Southern Ocean, Antarctic (Ross Sea)

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

F.E.Hoedt, J.H.Choat, J.J.Cruz, K.M.Neil & J.Collins, *Survey of the Port of Weipa: Port Marine Baseline Surveys and Surveys for Introduced Marine Pests*. Report by the School of Marine Biology, James Cook University, Townsville (2001)

A.McMinn & F.J.Scott, Dinoflagellates, *Antarctic Marine Protists* 202–250 (2005)