

**Gonyaulax verior** Sournia

*Beih. Nova Hedwigia* 48: 34 (1973)

*Amylax diacantha* Meunier *Mém. Mus. Roy. Hist. Nat. Belgique* 8(1): 74, pl. 19, figs 33–36 (1919); *Gonyaulax diacantha* (Meunier) J.Schiller, *Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz* 10(2), Teil 3(2): 300 (1935)

*Type locality*: North Sea (Belgium)

*Australian distribution*: N.S.W. (Port Hacking), Vic. (Port Phillip Bay; & cysts), Tas., S.A. (Port Lincoln; cysts)

*Extra-Australian distribution*: Mexico (Pacific coast), Gulf of Mexico, U.S.A. (Massachusetts), Canary Islands, British Isles, North Sea, Baltic Sea, Mediterranean Sea, Adriatic Sea, Black Sea, Taiwan, Japan

*References*: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)