

**Gymnodinium aureolum** (Hulburt) G.Hansen

in A.Hansen, N.Daugbjerg & P.Henriksen, *J. Phycol.* 36: 399, figs 22–36, 40, 44a, d–g, I (2000)

*Gyrodinium aureolum* Hulburt, *Biol. Bull.* 112: 209, pl. 2, figs 8, 9 (1957)

*Type locality:* U.S.A. (Massachusetts)

*Australian distribution:* Vic. (Gippsland Lakes), Tas. (Long Bay, Port Arthur, E coast; & cysts), S.A. (Adelaide)

*Extra-Australian distribution:* Argentina, U.S.A. (Virginia), Canary Islands, North Atlantic Ocean, British Isles, North Sea, Baltic Sea, Norwegian Sea, Mediterranean Sea, Adriatic Sea, South Korea, New Zealand

*Note:* An ichthyotoxic species

*References:* G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)