

Histioneis crateriformis F.Stein

Der Organismus der Infusionsthiere 3(2): 22, pl. 21, figs 5, 6 (1883)

Parahistioneis crateriformis (F.Stein) Kofoid & Skogsberg, *Mem. Mus. Comp. Zool. Harvard* 51: 590 (1928)

Type locality: Atlantic Ocean, "Südsee"

Histioneis reticulata Kofoid, *Bull. Mus. Compar. Zool. Harvard Coll.* 50: 205, pl. 15, fig. 95 (1907); *Parahistioneis reticulata* (Kofoid) Kofoid & Skogsberg, *Mem. Mus. Comp. Zool. Harvard* 51: 605 (1928)

Type locality: eastern tropical Pacific Ocean

Australian distribution: Coral Sea, N.S.W. (Trial Bay)

Extra-Australian distribution: Brazil, Gulf of Mexico, Atlantic Ocean, Canary Islands, Arabian Sea, Indian Ocean, NW Pacific Ocean, Solomon Islands, eastern tropical Pacific Ocean, SE Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

Synonymy after F.Gómez, A list of free-living dinoflagellate species in the world's oceans, *Acta Bot. Croat.* 64: 129–212 (2005)