

Histioneis milneri G.Murray & Whitting

Trans. Linn. Soc. London, Ser. Bot. 5: 334, pl. 33, fig. 1 ('1899') [1900]

Type locality: Atlantic Ocean

Histioneis helenae G.Murray & Whitting, *Trans. Linn. Soc. London*, Ser. Bot. 5: 335, pl. 33, fig. 2 ('1899') [1900]

Type locality: Atlantic Ocean

Histioneis hippoperoides Kofoid & Michener, *Bull. Mus. Comp. Zool. Harvard Coll.* 54(7): 296 (1911)

Type locality: eastern tropical Pacific Ocean

Australian distribution: NE Indian Ocean, southern Coral Sea, N.S.W. (Montague Island)

Extra-Australian distribution: Brazil, Gulf of Mexico, Atlantic Ocean, Canary Islands, Gulf of Aqaba, eastern tropical Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

E.J.F.Wood, Dinoflagellates in the Australian region. III – Further collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 17: 1–20 (1963)

Synonymy after F.Gómez, A list of free-living dinoflagellate species in the world's oceans, *Acta Bot. Croat.* 64: 129–212 (2005)