

**Karenia papilionacea** Haywood & Steid.

in A.J.Haywood, K.A.Steindinger, E.W.Truby, P.R.Bergquist, P.L.Bergquist, J.Adamson & L.Mackenzie, *J. Phycol.* 40: 167, figs 2e–h, 3a–f (2004)

*Type locality*: New Zealand

*Australian distribution*: Vic. (Gippsland Lakes), Tas. (Georges Bay, Moulting Bay, Ansons Bay, Little Swanport, Parsons Bay, D’Entrecasteaux Channe), S.A. (Port Lincoln), W.A. (Swan River)

*Extra-Australian distribution*: Gulf of Mexico, U.S.A. (Delaware), Spain (N coast), Mediterranean Sea, South China Sea, New Zealand

*References*: B.D.Mooney, M. de Salas, G.M.Hallegraeff & A.R.Price, Survey for karlotoxin production in 15 species of gymnodinioid dinoflagellates (Kareniaceae, Dinophyta), *J. Phycol.* 45: 164–175 (2009)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)