

Warnowia rosea (C.H.G.Pouchet) J.Schiller

Rabenhorst's *Kryptogamen-Flora von Deutschland, Österreich und der Schweiz* 10(3), Teil 2(4): 583, fig. 612 (1937)

Gymnodinium polyphemum var. *roseum* C.H.G.Pouchet, *J. Anat. Physiol.* 23: 96, pl. 10, fig. 1 (1877), as *polyphemus*

Type locality: France (Atlantic coast)

Australian distribution: Coral Sea, N.S.W., Vic., Tas.

Extra-Australian distribution: North America, British Isles, North Sea, Skaggerak-Kattegat, Norwegian Sea, Portugal, Mediterranean Sea, Indian Ocean, Japan, eastern Russia

References: E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)