

Oxytoxum caudatum J.Schiller

Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz. 10(3), Teil 2(3): 454, fig. 504 (1937)

Type locality: Adriatic Sea

Oxytoxum nanum Halldal, *Hvalrådets Skr.* 38: 56, fig. 20 (1953)

Type locality: Norwegian Sea

Australian distribution: Gulf of Carpentaria/North-west Shelf, N.S.W. (off Sydney)

Extra-Australian distribution: Brazil, Gulf of Mexico, North Atlantic Ocean, Norwegian Sea, Mediterranean Sea, Adriatic Sea, Black Sea, Indian Ocean, East China Sea, SW Pacific Ocean, Southern Ocean (S of Australia), Antarctica, Mexico (Pacific coast), SE Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

Synonymy after F.Gómez, A list of free-living dinoflagellate species in the world's oceans, *Acta Bot. Croat.* 64: 129–212 (2005)