

Oxytoxum crassum J.Schiller

Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz. 10(3), Teil 2(3): 459, fig. 518 (1937)

Type locality: Adriatic Sea

Oxytoxum globosum J.Schiller, *Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz.* 10(3), Teil 2(3): 458, fig. 515 (1937)

Type locality: Adriatic Sea

Australian distribution: East Australian Current, N.S.W. (off Sydney), E Indian Ocean

Extra-Australian distribution: Brazil, Gulf of Mexico, U.S.A. (Virginia), North Atlantic Ocean, British Isles, Mediterranean Sea, Adriatic Sea, Bay of Bengal, Southern Ocean (S of Australia), eastern tropical Pacific Ocean, SE Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

S.W.Jeffrey & G.M.Hallegraeff, Studies of phytoplankton species and photosynthetic pigments in a warm core eddy of the East Australian Current. I. Summer populations, *Mar. Ecol. Prog. Ser.* 3: 285–294 (1980)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

Synonymy after F.Gómez, A list of free-living dinoflagellate species in the world's oceans, *Acta Bot. Croat.* 64: 129–212 (2005)