

Oxytoxum sceptrum (F.Stein) B.Schröder

Mitt. Zool. Stat. Neapel 14: 18 (1900)

Pyrgidium sceptrum F.Stein, *Der Organismus der Infusionsthier* 3(2): pl. 5, figs 19–21 (1883)

Type locality: “Südsee”

Oxytoxum longiceps J.Schiller, *Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz* 10(3), Teil 2(3): 464, fig. 532 (1937)

Type locality: Adriatic Sea

Australian distribution: Coral Sea, Qld (Great Barrier Reef), East Australian Current, Tas. (off E coast), Indian Ocean W of Australia

Extra-Australian distribution: Brazil, Gulf of Mexico, Caribbean Sea, U.S.A. (Virginia), Canary Islands, Mediterranean Sea, Adriatic Sea, Aegean Sea, Black Sea, Indian Ocean, Bay of Bengal, South China Sea, East China Sea, Southern Ocean S of Australia (“cf.”), Tasman Sea, New Zealand, Pacific coast of Mexico, SE Pacific Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

N.Revelante, W.T.Williams & J.S.Bunt, Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef Australia, *Journal of Experimental Marine Biology and Ecology* 63: 27–46 (1982)