

Phalacroma circumsutum G.Karst.

Wiss. Ergeb. Deutschen Tiefsee-Exped. "Valdivia" 2(2): 421, pl. 53, fig. 8 (1907)

Dinophysis circumsuta (G.Karst.) Balech, *Revta Mus. Argent. Cienc. Nat. "Bernardino Rivadavia" (Hidrobiol.) 2: 82 (1967)*

Type locality: Indian Ocean

Australian distribution: Coral Sea, N.S.W. (off Sydney), East Australian Current, North-west Shelf

Extra-Australian distribution: Argentina, Brazil, Gulf of Mexico, Indian Ocean (Zanzibar), South China Sea, Antarctica

References: E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO 14: 1–55 (1963)*

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser. 20: 59–74 (1984)*

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res. 37: 361–377 (1986)*

G.M.Hallegraeff & I.A.N.Lucas, The marine dinoflagellate genus *Dinophysis* (Dinophyceae): photosynthetic, neritic and non-photosynthetic, oceanic species, *Phycologia 27: 25–42 (1988)*