

**Phalacroma doryphorum** F.Stein

*Der Organismus der Infusionsthiere* 3(2): 23, pl. 19, figs 1–4 (1883)

*Dinophysis doryphora* (F.Stein) T.H.Abé, *Publ. Seto Mar. Biol. Lab.* 15(1): 77 (1967)

Type locality: Atlantic Ocean, “Südsee”

Australian distribution: Coral Sea, East Australian Current, N.S.W. (off Sydney), North-west Shelf

Extra-Australian distribution: Brazil, Gulf of Mexico, Atlantic Ocean, Canary Islands, Mediterranean Sea, Indian Ocean, Mexico (Pacific coast)

References: G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

G.M.Hallegraeff & I.A.N.Lucas, The marine dinoflagellate genus *Dinophysis* (Dinophyceae): photosynthetic, neritic and non-photosynthetic, oceanic species, *Phycologia* 27: 25–42 (1988)