

**Phalacroma parvulum** (F.Schütt) Jörg.

*Rep. Danish Ocean. Exped., 1908–10, Mediterr. Seas II, J, 2: 7 (1923)*

*Phalacroma porodictyum* var. *parvula* F.Schütt, *Ergebn. Plankton-Exped. Humboldt-Stiftung* 4: 148, pl. 2, fig. 13 (1895); *Dinophysis parvula* (F.Schütt) Balech, *Revta Mus. Argent. Cienc. Nat. “Bernardino Rivadavia” (Hidrobiol.)* 2: 83 (1967)

Type locality: Atlantic Ocean

Australian distribution: Coral Sea, East Australian Current, N.S.W. (off Sydney), Tas., North-west Shelf

Extra-Australian distribution: Brazil, U.S.A. (Virginia), Atlantic Ocean, British Isles, Mediterranean Sea, Black Sea, Indian Ocean

References: E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

G.M.Hallegraeff & I.A.N.Lucas, The marine dinoflagellate genus *Dinophysis* (Dinophyceae): photosynthetic, neritic and non-photosynthetic, oceanic species, *Phycologia* 27: 25–42 (1988)