

**Podolampas elegans** F.Schütt

*Ergebn. Plankton-Exped. Humboldt-Stiftung* 4: 53, 91, pl. 18, fig. 57 (1895)

*Type locality:* Atlantic Ocean

*Australian distribution:* Gulf of Carpentaria/North-west Shelf, Coral Sea, N.S.W. (off Sydney), East Australian Current

*Extra-Australian distribution:* Brazil, Gulf of Mexico, U.S.A. (Virginia), Canary Islands, Atlantic Ocean, Mediterranean Sea, Adriatic Sea, Black Sea, Bay of Bengal, Indian Ocean, Philippines, South China Sea, East China Sea, tropical central Pacific Ocean, Mexico (Gulf of California), Peru-Chile Current

*References:* E.J.F.Wood, Dinoflagellates in the Australian region. II – Recent collections, *Tech. Pap. Div. Fish. Oceanogr. CSIRO* 14: 1–55 (1963)

S.W.Jeffrey & G.M.Hallegraeff, Studies of phytoplankton species and photosynthetic pigments in a warm core eddy of the East Australian Current. I. Summer populations, *Mar. Ecol. Prog. Ser.* 3: 285–294 (1980)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)