

Pronoctiluca acuta (Lohmann) J.Schiller

Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz 10(2), Teil 3(1): 271 (1932)

Rhynchomonas acuta Lohmann, *Veröff. Inst. Meeresk. Univ. Berlin. Neue Folge. A, Geogr.-Naturwiss. Reihe* 1: 24, text fig. 2 (1912)

Type locality: Atlantic Ocean [Brazilian waters]

Rhynchomonas curvata Lohmann, *Int. Rev. Hydrobiol. Hydrogeogr.* 5: 362, fig. 17.3 (1913); *Pronoctiluca acuta* var. *curvata* (Lohmann) J.Schiller, *Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz* 10(2), Teil 3(1): 272 (1932)

Type locality: SW Atlantic Ocean

Australian distribution: N.S.W. (Sydney Harbour)

Extra-Australian distribution: SW Atlantic Ocean, Brazil, Gulf of Mexico, Canary Islands, Mediterranean Sea, Adriatic Sea, Black Sea, Bay of Bengal, New Caledonia, equatorial Pacific Ocean

Note: Marine and estuarine

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)