

**Protoperidinium avellana** (Meunier) Balech

*Revta Mus. Argent. Cienc. Nat. "Bernardino Rivadavia" (Hidrobiol.)* 4: 54 (1974)

*Properidinium avellana* Meunier, *Mém. Mus. Roy. Hist. Nat. Belgique* 8(1): 546, pl. 18, figs 37–41 (1919)

Type locality: North Sea (Belgium)

Australian distribution: N.S.W. (off Sydney), Vic. (Port Phillip Bay; cysts), Tas. (cysts)

Extra-Australian distribution: Brazil, Mexico (Pacific coast), Gulf of Mexico, U.S.A. (Virginia), British Isles, North Sea (Belgium), Mediterranean Sea, Arctic Ocean, far-eastern Russia, East China Sea, South China Sea, Coral Sea, Chile

References: G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)