

Protoperidinium bipes (Paulsen) Balech

Revta Mus. Argent. Cienc. Nat. "Bernardino Rivadavia" (Hidrobiol.) 4: 53 (1974)

Glenodinium bipes Paulsen, *Medd. Kommn. Havunders. Kobenh. Ser. Plankton* 1: 21, figs 3, 4 (1904)

Type locality: Iceland

Peridinium minusculum Pavill., *Trav. Inst. Bot. Univ. Montpellier* 2: 57, pl. 3, figs 7–9 (1905)

Type locality: Mediterranean Sea (France)

Australian distribution: East Australian Current, N.S.W. (Sydney)

Extra-Australian distribution: U.S.A. (Maine), Canada (Bay of Fundy), Canary Islands, Iceland, Kara Sea, Svalbard, British Isles, North Sea, Baltic Sea, Mediterranean Sea (France), Black Sea, Japan, South Korea

Note: Marine and estuarine

References: S.W.Jeffrey & G.M.Hallegraeff, Studies of phytoplankton species and photosynthetic pigments in a warm core eddy of the East Australian Current. I. Summer populations, *Mar. Ecol. Prog. Ser.* 3: 285–294 (1980)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)