

Protoperidinium conicoides (Paulsen) Balech

Revta Mus. Argent. Cienc. Nat. "Bernardino Rivadavia" (Hidrobiol.) 3: 356, pl. 3, fig. 2 (1973)

Peridinium conicoides Paulsen, *Meddel. Komm. Havundersøg., Ser. Plankton* 1(3): 3, fig. 2 (1905)

Type locality: ?

Australian distribution: N.S.W. (Port Hacking), Vic. (cysts), Tas. (cysts), W.A. (Albany)

Extra-Australian distribution: Gulf of Mexico, U.S.A. (Massachusetts), North Atlantic Ocean, Arctic Ocean, British Isles, Baltic Sea, North Sea, Black Sea, Indian (S coast), Japan, Antarctica, Chile, Peru

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

C.J.Bolch & G.M.Hallegraeff, Dinoflagellate cysts from Recent marine sediments of Tasmania, Australia, *Bot. Mar.* 33: 173–192 (1990)

J.A.Sonneman & D.R.A.Hill, A taxonomic survey of cyst-producing dino-flagellates from the coastal waters of Victoria, Australia, *Bot. Mar.* 40: 149–177 (1997)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)