

**Protoperidinium divergens** (Ehrenb.) Balech

*Revta Mus. Argent. Cienc. Nat. “Bernardino Rivadavia” (Hidrobiol.)* 4: 60 (1974)

*Peridinium divergens* Ehrenb., *Deutsche Akad. Wiss. Berlin* 1840: 201 (1840)

Type locality: Baltic Sea

Australian distribution: Gulf of Carpentaria/North-west Shelf, Qld to Tas. (Maria Island), S.A., W.A. (Albany)

Extra-Australian distribution: Brazil, Canary Islands, British Isles, North Sea, Baltic Sea, Black Sea, Red Sea, Mexico (Pacific coast), U.S.A. (Washington)

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)