

Protoperidinium globulum (F.Stein) Balech

Revta Mus. Argent. Cienc. Nat. "Bernardino Rivadavia" (Hidrobiol.) 4: 64 (1974)

Peridinium globulus F.Stein, *Organismus der Infusionstiere* 3(2): pl. 9, figs 5–7 (1883)

Type locality: Mediterranean Sea, Atlantic Ocean, "Südsee"

Australian distribution: E coast, N.S.W. (Sydney), Vic. (Port Fairy), Tas. (Bass Str., River Derwent), W.A. (Fremantle)

Extra-Australian distribution: North Atlantic Ocean, Canary Islands, British Isles, North Sea, Svalbard, Mediterranean Sea, Adriatic Sea, Black Sea, Indian Ocean, Pacific Ocean

Note: Marine and estuarine

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)