

Protoperidinium oblongum (Auriv.) Parke & J.D.Dodge

in Parke & Dixon, *J. Mar. Biol. Assoc. U.K.* 56: 545 (1976)

Peridinium divergens var. *oblongum* Auriv., *Kungl. Svenska Vetensk.-Akad. Handl.* 30(3): 96 (1898)

Type locality: Great Britain

Australian distribution: Gulf of Carpentaria/North-west Shelf, Qld, N.S.W., Vic. (Port Phillip Bay; & cysts), Tas., W.A. (Albany)

Extra-Australian distribution: Gulf of Mexico, Caribbean Sea, Canada (Gulf of St. Lawrence), Canary Islands, Spain, North Atlantic Ocean, British Isles, North Sea, Baltic Sea, Mediterranean Sea, Black Sea, Persian Gulf, Arabian Sea, Papua New Guinea, Solomon Islands, SW Pacific Ocean, Heard Island, eastern Pacific Ocean (Chile-Peru Current)

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)

C.J.Bolch & G.M.Hallegraeff, Dinoflagellate cysts from Recent marine sediments of Tasmania, Australia, *Bot. Mar.* 33: 173–192 (1990)

J.A.Sonneman & D.R.A.Hill, A taxonomic survey of cyst-producing dino-flagellates from the coastal waters of Victoria, Australia, *Bot. Mar.* 40: 149–177 (1997)

A.McMinn, C.J.S.Bolch, M.F. de Salas & G.M.Hallegraeff, Recent dinoflagellate cysts, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 260–292 (2010)