

Protoperidinium pallidum (Ostenf.) Balech

Revta Mus. Argent. Cienc. Nat. "Bernardino Rivadavia" (Hidrobiol.) 3: 365 (1973)

Peridinium pallidum Ostenf., in M.Knudsen & C.[H.]Ostenfeld, *Iagttagelser over overfladevants Temperature, Saltoldighet og Plankton paa Islandske og Grönlandske Skibsrouter I* 1898 60 (1899)

Type locality: North Atlantic Ocean (Denmark Strait)

Australian distribution: Gulf of Carpentaria/North-west Shelf, N.S.W. (off Sydney); “rare — Port Hacking, N.S.W., to Queensland” [Wood, 1954]

Extra-Australian distribution: Canary Islands, North Atlantic Ocean (Denmark Strait), British Isles, Norway, Adriatic Sea, Black Sea, West Africa, Japan

References: E.J.F.Wood, Dinoflagellates in the Australian region, *Austral. J. Mar. Freshw. Res.* 5: 171–351 (1954)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff & D.D.Reid, Phytoplankton species successions and their hydrological environment at a coastal station off Sydney, *Austral. J. Mar. Freshw. Res.* 37: 361–377 (1986)