

Takayama helix de Salas, Bolch, Botes & Hallegr.

in M.F. de Salas, C.J.S.Bolch, L.Botes, G.Nash, S.W.Wright & G.M.Hallegraeff, *J. Phycol.* 39: 1238, figs 14–28, 34, 35 (2003)

Type locality: Australia (Tas.)

Australian distribution: Vic. (Gippsland Lakes), Tas. (E coast), S.A. (Port Lincoln)

Extra-Australian distribution: Spain, South Africa, Japan, New Zealand

References: M.F. de Salas, C.J.S.Bolch, L.Botes, G.Nash, S.W.Wright & G.M.Hallegraeff, *Takayama* (Gymnodiniales, Dinophyceae) gen. nov., a new genus of unarmoured dinoflagellates with sigmoid apical grooves, including the description of two new species, *J. Phycol.* 39: 1233–1246 (2003)

B.D.Mooney, M. de Salas, G.M.Hallegraeff & A.R.Price, Survey for karlotoxin production in 15 species of gymnodinoid dinoflagellates (Kareniaceae, Dinophyta), *J. Phycol.* 45: 164–175 (2009)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)