

**Takayama tasmanica** de Salas, Bolch & Hallegr.

in M.F. de Salas, C.J.S.Bolch, L.Botes, G.Nash, S.W.Wright & G.M.Hallegraeff, *J. Phycol.* 39: 1235, figs 2–12, 32, 33 (2003)

*Type locality:* Australia (Tas.)

*Australian distribution:* Tas. (River Derwent, E coast)

*Extra-Australian distribution:* Gulf of Mexico, New Zealand

*References:* M.F. de Salas, C.J.S.Bolch, L.Botes, G.Nash, S.W.Wright & G.M.Hallegraeff, *Takayama* (Gymnodiniales, Dinophyceae) gen. nov., a new genus of unarmoured dinoflagellates with sigmoid apical grooves, including the description of two new species, *J. Phycol.* 39: 1233–1246 (2003)

B.D.Mooney, M. de Salas, G.M.Hallegraeff & A.R.Price, Survey for karlotoxin production in 15 species of gymnodinioid dinoflagellates (Kareniaceae, Dinophyta), *J. Phycol.* 45: 164–175 (2009)

G.M.Hallegraeff, C.J.S.Bolch, J.M.Huisman & M.F. de Salas, Planktonic dinoflagellates, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 145–212 (2010)