

Azpeitia nodulifera (A.F.W.Schmidt) Fryxell & P.A.Sims

in Fryxell *et al.*, *Syst. Bot. Monogr.* 13: 19, fig. 17(1A–3), fig. 18(1, 2, 4, 5), fig. 30 (3, 4) (1986)

Coscinodiscus nodulifera A.F.W.Schmidt, in Schmidt *et al.*, *Atlas der Diatomaceen-kunde*, Ser. 2, 15/16: pl. 59, figs 20–23 (1878), as *nodulifer*

Type locality: U.S.A. (California, Florida)

Australian distribution: N & NW Australia

References: G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

F.J.Scott & D.P.Thomas, Diatoms, *Antarctic Marine Protists* 13–201 (2005)

Extra-Australian distribution: Gulf of California, Gulf of Mexico, U.S.A. (California, Florida), Spain, British Isles, North Sea, Baltic Sea, Black Sea, Gulf of Suez, SE Atlantic Ocean (Benguela Current), Gulf of Thailand, South China Sea, Japan, Sea of Okhotsk, Southern Ocean (S of Australia), East Antarctica