

Cerataulina pelagica (Cleve) Hendey

Discovery Rep. 16: 279 (1937)

Zygoceros pelagica Cleve, *Det Videnskabelige Udbytte af Kanonbaaden "Hauchs" Togter i de Danske Have Indenfor Skagen I Aarene 1883–86*: 54 (1889)

Type locality: Denmark

Australian distribution: N & NW Australia, Qld (SE coast), N.S.W. (E and SE coasts), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 49 (2010)

Extra-Australian distribution: Gulf of Mexico, Canada (Nova Scotia), Canary Islands, British Isles, North Sea, Spain, Black Sea, Japan