

Chaetoceros curvisetus Cleve

Det Videnskabelige Udbytte af Kanonbaaden "Hauchs" Togter i de Danske Have Indenfor Skagen I Aarene 1883–86: 55 (1889)

Type locality: Denmark

Australian distribution: Qld (Moreton Bay), N.S.W., Vic., Tas. (incl. Bass. Str.), W.A. (Shark Bay).

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 39 (2010)

Extra-Australian distribution: Argentina, Gulf of Mexico, U.S.A. (Washington, Alaska, NE coast), Canada (New Brunswick, Nova Scotia), British Isles, North Sea, Arctic Ocean, Canary Islands, Baltic Sea, Spain, Portugal, Black Sea, Sea of Japan, China, W Pacific Ocean, Pacific coast of South America, Southern Ocean, Antarctica