

Cocconeis scutellum Ehrenb. var. **scutellum**

Die Infusionsthierchen als vollkommene Organismen 194, pl. 14, fig. 8 (1838)

Type locality: Sweden, Germany

Australian distribution: N.T., Qld, N.S.W. (Port Hacking, Lake Conjola), Vic. (Gippsland Lakes, Port Phillip Bay, Western Port Bay), Tas. (Ansons Bay, Georges Bay, Orielton Lagoon, Pipeclay Lagoon), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally epontic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

N.Foged, Diatoms in eastern Australia, *Biblioth. Phycol.* 41: 1–243 (1978)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

K.Saunders, C.Lane, S.Cook, A.McMinn & G.M.Hallegraeff, Benthic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 87 (2010)

Extra-Australian distribution: Canada (E coast), Canary Islands, North Sea, Baltic Sea, Mediterranean Sea, Black Sea, Japan, New Zealand, Hawaiian Islands, Antarctica

Note: Estuarine and in fresh water