

Diploneis smithii (Bréb.) Cleve var. **smithii**

Kongl. Svenska Vetensk.-Akad. Handl., ser. 2, 26: 96 (1894)

Navicula smithii Bréb., in Smith, *Syn. Brit. Diatomaceae* 2: 92 (1856)

Type locality: Great Britain

Australian distribution: Qld, N.S.W. (Lake Macquarie, Botany Bay, Port Hacking), Vic., Tas., W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally epontic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

N.Foged, Diatoms in eastern Australia, *Biblioth. Phycol.* 41: 1–243 (1978)

K.Saunders, C.Lane, S.Cook, A.McMinn & G.M.Hallegraeff, Benthic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 108 (2010)

Extra-Australian distribution: U.S.A. (Florida, Virginia), Canada (E coast), Canary Islands, British Isles, North Sea, Baltic Sea, Spain, Egypt, Black Sea, Persian Gulf, W India, Sri Lanka, Japan, New Zealand, Antarctica, near North Pole