

Grammatophora marina (Lyngb.) Kütz.

Kieselschaligen Bacillarien Diatomeen 128, pl. 17, fig. 24 (1844)

Diatoma marinum Lyngb., *Tent. Hydrophytol. Danicae* 180 (1819)

Type locality: Baltic Sea

Australian distribution: Qld (Moreton Bay), N.S.W. (Port Stephens, Port Jackson, Port Hacking, Botany Bay, Lake Macquarie), Vic. (Gippsland Lakes, Port Phillip Bay, Western Port Bay), Tas. (Georges Bay), W.A. (King George Sound)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally eponitic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

N.Foged, Diatoms in eastern Australia, *Biblioth. Phycol.* 41: 1–243 (1978)

K.Saunders, C.Lane, S.Cook, A.McMinn & G.M.Hallegraeff, Benthic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 123 (2010)

Extra-Australian distribution: U.S.A. (NE coast), Canada (Nova Scotia), Canary Islands, Baltic Sea, Spain, Romania, South Africa, Japan, Pacific Ocean, New Zealand, Campbell Island, Southern Ocean