

Halamphora coffeaeformis (C.Agardh) Levkov

Diatoms of Europe 5: 179, pl. 91, figs 1–14; pl. 94, figs 17–27; pl. 99, figs 15–23 (2009)

Frustulia coffeaeformis C.Agardh, *Flora* 10: 627 (1827)

Amphora coffeaeformis (C.Agardh) Kütz., *Kieselschaligen Bacillarien Diatomeen* 108, pl. 5, fig. 37 (1844)

Type locality: Sweden

Australian distribution: N.T., Qld, N.S.W., S.A., W.A. (Perth)

References: E.J.F.Wood, Studies on Australian and New Zealand diatoms. IV. Descriptions of further sedentary species, *Trans. & Proc. Roy. Soc. New Zealand* 88: 669–698 (1961)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

K.Saunders, C.Lane, S.Cook, A.McMinn & G.M.Hallegraeff, Benthic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 126 (2010)

Extra-Australian distribution: Argentina, U.S.A. (Oklahoma), Europe, Kara Sea, Canary Islands, Spain, France, Sweden, Czech Republic, Israel, Iran, Hawaiian Islands, Antarctica

Note: In brackish water and marine and hypersaline habitats