

Lyrella lyra (Ehrenb.) Karajeva var. **lyra**

Bot. Zhurn. 63: 1595 (1978)

Navicula lyra Ehrenb., *Abh. Königl. Akad. Wiss. Berlin* 1841: 131 (1843)

Type locality: Falkland Islands

Australian distribution: Qld (Noosa), N.S.W. (Port Hacking), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally epontic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

N.Foged, Diatoms in eastern Australia, *Biblioth. Phycol.* 41: 1–243 (1978), as var. *typica*

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

Extra-Australian distribution: Falkland Islands, S Brazil, Mexico (Baja California), Gulf of Mexico, Caribbean Sea, U.S.A. (Washington, Great Lakes, South Carolina, Maine), Canada (New Brunswick, Nova Scotia), Canary Islands, British Isles, North Sea, Baltic Sea, Adriatic Sea, Black Sea, Arabian Sea, Sri Lanka, Bay of Bengal, W Pacific Ocean, Hawaiian Islands

Note: Usually in brackish water