

Navicula tripunctata (O.F.Müll.) Bory var. **tripunctata**

Dict. Class. Hist. Nat. 2: 128 (1822)

Vibrio tripunctatus O.F.Müll., *Anim. Infus. Fluv. & Mar.* 52, pl. 7, fig. 2a, b (1786)

Type locality: Denmark

Navicula gracilis Ehrenb., *Abh. Königl. Akad. Wiss. Berlin* 1832: 64, 68 (1832)

Type locality: ?

Australian distribution: Qld, N.S.W. (Lake Conjola), Vic., Tas., W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally eponic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

K.Saunders, C.Lane, S.Cook, A.McMinn & G.M.Hallegraeff, Benthic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 113 (2010)

Extra-Australian distribution: U.S.A. (E coast), Canada (New Brunswick), British Isles, North Sea, Baltic Sea, E Europe, Spain, Romania, Turkey, Iran, South Africa, China

Note: Usually in fresh water; also estuarine and in other brackish habitats