

Plagiogramma staurophorum (W.Greg.) Heib.

Consp. Crit. Diatom. Dan. 55 (1863)

Denticula staurophora W.Greg., *Trans. Roy. Soc. Edinburgh* 21: 495, pl. 2, fig. 37 (1857)

Type locality: Great Britain

Australian distribution: Qld (Heron Island), N.S.W. (Byron Bay, Port Hacking), Vic. (Lakes Entrance), Tas., W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally epontic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

N.Foged, Diatoms in eastern Australia, *Biblioth. Phycol.* 41: 1–243 (1978)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

K.Saunders, C.Lane, S.Cook, A.McMinn & G.M.Hallegraeff, Benthic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 137 (2010)

Extra-Australian distribution: Argentina, Brazil, Gulf of Mexico, U.S.A. (E coast), Arctic Canada, British Isles, North Sea, Baltic Sea, France, Spain