

**Podosira stelligera** (Bailey) A.Mann.

*Contrib. U.S. Natl Herb.* 10: 242 (1907)

*Hyalodiscus stelligera* Bailey, *Smithsonian Contrib. Knowl.* 7(3): 10 (1854)

*Type locality:* U.S.A. (Florida)

*Australian distribution:* N & NW Australia, Qld (Capricorn Coast, Noosa), N.S.W. (South Coast), Vic. (Lakes Entrance), W.A. (Perth)

*References:* L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

N.Foged, Diatoms in eastern Australia, *Biblioth. Phycol.* 41: 1–243 (1978)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

*Extra-Australian distribution:* Gulf of Mexico, U.S.A. (Florida), British Isles, France, Spain, North Sea, Baltic Sea, Yellow Sea, western Pacific Ocean, New Zealand