

**Proboscia indica** (H.Perag.) Hernández-Becerril

*Diatom Res.* 10: 254, figs 5, 6 (1995)

*Rhizosolenia indica* H.Perag., *Diatomiste* 1(9): 116, pl. 5, fig. 16; pl. 18, fig. 16 (1892)

*Rhizosolenia alata* f. *indica* Ostenf., in Ostenfeld & Schmidt, *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1901: 160 (1901)

*Type locality:* Indian Ocean

*Australian distribution:* N & NW Australia, N.S.W. (Port Hacking)

*References:* L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

*Extra-Australian distribution:* Argentina, S Brazil, Mexico (Pacific coast), Venezuela, U.S.A. (California, NE coast), Bay of Biscay, British Isles, North Atlantic Ocean, North Sea, Baltic Sea, Mediterranean Sea, Adriatic Sea, Red Sea, Persian Gulf, Indian Ocean, SE Pacific Ocean, New Zealand