

Pseudonitzschia americana (Hasle) Fryxell

in Hasle, *Beih. Nova Hedwigia* 106: 318 (1993)

Nitzschia americana Hasle, *Skr. Norske Vidensk.-Akad. Oslo, Math.-Naturvidensk. Kl., n. ser.*, 16: 41, pl. 1, fig. 4; pl. 14, figs 13–19 (1964)

Type locality: Uruguay

Australian distribution: N.T./Qld (Gulf of Carpentaria), Tas. (River Derwent)

References: G.M.Hallegraeff, Species of the diatom genus *Pseudo-nitzschia* in Australian waters, *Bot. Mar.* 37: 397–411 (1994)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 66 (2010)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

Extra-Australian distribution: Uruguay, Chile, Peru, Pacific coast of Mexico, Gulf of Mexico, U.S.A. (California, NE coast), Canada (Bay of Fundy), Netherlands, South Africa, Vietnam, Japan