

Pseudosolenia calcaravis (M.Schultze) Sundström

Mar. Diatom Genus Rhizosolenia 95, figs 40–46, 247–257 (1986)

Rhizosolenia calcaravis M.Schultze, *Arch. Anat. Physiol. Wiss. Med.* 1858: 339, pl. 13, figs 5–10 (1858)

Type locality: ?

Australian distribution: N & NW Australia, Coral Sea, Qld (Thursday Island to S Coast), S.A. (Gulf St. Vincent), W.A. (Perth to Scotts Reef)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

Extra-Australian distribution: Argentina, Gulf of Mexico, Caribbean Sea, U.S.A. (NE coast), Canada (Nova Scotia), Bay of Biscay, Spain, Portugal, British Isles, North Atlantic Ocean, Mediterranean Sea (Spain, France, Turkey), Adriatic Sea, Black Sea, Caspian Sea, Sea of Azov, Gulf of Guinea, Persian Gulf, Bay of Bengal, Yellow Sea, Sea of Japan, East China Sea, Papua New Guinea, Solomon Islands, Vanuatu, New Caledonia, Fiji, Tasman Sea, W Pacific Ocean, New Zealand