

Shionodiscus oestrupii var. **oestrupii** (Ostenf.) A.J.Alverson, S.H.Kang & E.C.Theriot

Diatom Res. 21: 258 (2006)

Coscinosira oestrupii Ostenf., *Iagttagelser Overfladevandets Temkperature, Saltholdighed og Plankton paa islandske og gronlandske Skibsrouter i 1899* 52 (1900)

Thalassiosira oestrupii (Ostenf.) Hasle, *Taxon* 21: 544, figs 1–10 (1972)

Type locality: Greenland/Iceland?

Australian distribution: S of Tas.

References: F.J.Scott & D.P.Thomas, *Diatoms, Antarctic Marine Protists* 13–201 (2005)

O.E.Romero, L.K.Armand, X.Crosta, J.-J.Pichon, The biogeography of major diatom taxa in Southern Ocean surface sediments: 3. Tropical/subtropical species, *Palaeogeography, Palaeoclimatology, Palaeoecology* 223: 49–65 (2005)

Extra-Australian distribution: South Atlantic Ocean, Gulf of Mexico, U.S.A. (Gulf of Maine), Canada (Bay of Fundy), Greenland/Iceland (?), North Atlantic Ocean, British Isles, Norwegian Sea, Black Sea, Persian Gulf, Indian Ocean, Sea of Japan, Yellow Sea, Southern Ocean, Tasman Sea, New Zealand, Antarctica, Pacific coast of South America