

Stephanopyxis turris (Grev.) Ralfs

in Pritchard, *History of the Infusoria, including the Desmidiaceae and Diatomaceae, British and Foreign* 826, pl. 4, fig. 74 (1861)

Cresswellia turris Grev., in Gregory, *Trans. Roy. Soc. Edinburgh* 21: 538, pl. 6, fig. 109 (1857)

Type locality: Great Britain

Australian distribution: Qld (Torres Str.), N.S.W. (Port Stephens, Sydney), Vic., Tas. (Bass. Str.), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 19 (2010)

Extra-Australian distribution: Canada (Bay of Fundy), British Isles, North Sea, Mediterranean Sea, Canary Islands, Black Sea, SW Asia, New Zealand, Pacific coast of South America

Note: A marine and estuarine species