

Terpsinoe americana (Bailey) Ralfs

in Pritchard, *History of the Infusoria, including the Desmidiaceae and Diatomaceae, British and Foreign* 859 (1861)

Tetragramma americana Bailey, *Smithsonian Contrib. Knowl.* 7(3): 7, figs 1, 2 (1854)

Type locality: U.S.A. (Maine)

Australian distribution: Qld (Cairns), N.S.W. (Port Hacking), Vic. (Anderson Inlet), Tas. (Little Swanport), W.A. (Perth)

References: E.J.F.Wood, Studies on Australian and New Zealand diatoms. IV. Descriptions of further sedentary species, *Trans. & Proc. Roy. Soc. New Zealand* 88: 669–698 (1961)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

K.Saunders, C.Lane, S.Cook, A.McMinn & G.M.Hallegraeff, Benthic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 135 (2010)

Extra-Australian distribution: Uruguay, Brazil, Pacific coast of Mexico, Gulf of Mexico, Venezuela, U.S.A. (NE coast), North Sea, Baltic Sea, Spain, E Mediterranean Sea, Russia (Sea of Okhotsk), Japan, New Zealand