

Thalassionema nitzschioides (Grunow) Mereschk. var. **nitzschioides**

Scr. Bot. Horti Univ. Petropol. 19: 78 (1902)

Synedra nitzschioides Grunow, *Verhandl. Kaiserl.-Königl. Zool.-Bot. Ges. Wien* 12: 403, pl. 5, fig. 18a, b (1862)

Type locality: ?

Australian distribution: N.T./Qld (Gulf of Carpentaria), Qld (Cape York, E coast), N.S.W. (North Coast, Central Coast, South Coast), Tas., W.A. (Perth, Rottnest Island, NW Shelf)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. II. Normally eponic and benthic genera, *Trans. & Proc. Roy. Soc. New Zealand* 86: 1–58 (1959)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

G.M.Hallegraeff, Taxonomy and morphology of the marine plankton diatoms *Thalassionema* and *Thalassiothrix*, *Diatom Res.* 1: 57–80 (1986)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 74 (2010) N.Revelante & M.Gilmartin, Characteristics of the microplankton and nanoplankton communities of an Australian coastal plain estuary, *Austral. J. Mar. Freshw. Res.* 29: 9–18 (1978)

Extra-Australian distribution: Argentina, South Atlantic Ocean, North Atlantic Ocean, North Sea, Baltic Sea, Arctic Ocean, Black Sea, Mediterranean Sea, Indian Ocean, Indonesia, Timor Sea, Japan, Korean Peninsula, W Pacific Ocean, N Pacific Ocean, New Zealand, Southern Ocean, Antarctica