

Thalassiosira eccentrica (Ehrenb.) Cleve

Conseil Permanent pour l'Exploration de la Mer, Bulletin des Résultats aquit pendant les course périodiques 1903–1904: 216 (1904)

Coscinodiscus eccentricus Ehrenb., *Abh. Königl. Akad. Wiss. Berlin* 1839: 146 (1841), as *excentricus*

Type locality: Mediterranean Sea (Algeria; fossil), North Sea (Germany), Kattegat, Mexico (living)

Australian distribution: N.T./Qld (Gulf of Carpentaria), Qld (Capricorn Coast), East Australian Current, N.S.W. (Lake Macquarie), Vic. (Lakes Entrance, Port Phillip Bay), Great Australian Bight, W.A. (Perth, NW Shelf)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

N.Foged, Diatoms in eastern Australia, *Biblioth. Phycol.* 41: 1–243 (1978)

J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

G.M.Hallegraeff, Species of the diatom genus *Thalassiosira* in Australian waters, *Bot. Mar.* 27: 495–513 (1984)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

I.Jameson & G.M.Hallegraeff, Planktonic diatoms, *Algae of Australia: Phytoplankton of Temperate Coastal Waters* 59 (2010)

Extra-Australian distribution: Chile, Peru, Mexico, U.S.A. (Washington, E coast), British Isles, North Sea, Baltic Sea, Bay of Biscay, Spain, Canary Islands, Arabian Sea, Indian Ocean, Japan, China, Korean Peninsula, New Zealand, Campbell Island, Norfolk Island, Southern Ocean, Antarctica