

Thalassiosira subtilis (Ostenf.) Gran

Nytt Mag. Naturvidensk. 38: 117 (1900)

Podosira subtilis Ostenf., *Iagttagelser Overfladevandets Temkperature, Saltholdighed og Plankton paa islandske og gronlandske Skibsrouter i 1898* 55 (1899)

Type locality: North Atlantic Ocean

Australian distribution: N.T./Qld (Gulf of Carpentaria), Qld (SE coast), N.S.W. (Central Coast), Vic., Tas. (Bass Str., E coast), East Australian Current, W.A. (NW Shelf)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

G.M.Hallegraeff, Species of the diatom genus *Thalassiosira* in Australian waters, *Bot. Mar.* 27: 495–513 (1984)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

Extra-Australian distribution: North Atlantic Ocean, British Isles, North Sea, Spain, Portugal, China, Japan, W Pacific Ocean, North Pacific Ocean, New Zealand