

**Tryblionella apiculata** W.Greg.

*Trans. Microscop. Soc. London* 5: 79, pl. 1, fig. 43 (1857)

*Nitzschia apiculata* (W.Greg.) Grunow, *Naturwissenschaftliche Beiträge zur Kenntnis der Kaukasusländer, auf Grund seiner Sammelbeute* 118 (1878)

Type locality: Great Britain

*Synedra constricta* Kütz., *Kieselschaligen Bacillarien Diatomeen* 64, pl. 3, fig. 70 (1844)

*Nitzschia constricta* (Kütz.) Ralfs, in Pritchard, *History of the Infusoria, including the Desmidiaceae and Diatomaceae, British and Foreign* 780 (1861)

*Tryblionella constricta* (Kütz.) Poulin, in Poulin et al., *Naturaliste Canadien* 117: 96; figs 107, 108 (1990), nom. illeg., non *Tryblionella constricta* W.Greg. (1855)

Type locality: Adriatic Sea (Italy)

Australian distribution: N & NW Australia, W.A. (Perth)

References: J.John, The diatom flora of the Swan River Estuary, Western Australia, *Biblioth. Phycol.* 64: 1–358 (1983)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

Extra-Australian distribution: U.S.A., Canada (Quebec), Canary Islands, Spain, British Isles, Czech Republic, Adriatic Sea (Italy), Caspian Sea

Note: In brackish and freshwater habitats